

Accuration® Balancer 26 R300

**Specie:** Beef**Type of Feed:** Supplement**Form of Feed:** Pellet**General Description:**

A supplement for mixing with coarse cracked corn or shelled corn and by-product ingredients for finishing cattle weighing over 650 lbs. or dairy beef steers weighing over 400 lbs. that is formulated to provide intake modifying properties to the complete ration.

PURINA®

ACCURATION® BALANCER 26 R300

TYPE B MEDICATED FEED

SUPPLEMENT FEED FOR GROWER-FINISHER FEEDLOT CATTLE

For improved feed efficiency in cattle fed in confinement for slaughter.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 300.00 g/ton

GUARANTEED ANALYSIS

Crude Protein, (Min) 26.00 %
 (This includes not more than 17.00% equivalent crude protein from non-protein nitrogen.)

Crude Fat, (Min) 2.00 %

Crude Fiber, (Max) 15.00 %

Calcium (Ca), (Min) 6.50 %

Phosphorus (P), (Min) 0.50 %

Salt (NaCl), (Min) 2.50 %

Potassium (K), (Min) 2.00 %

Copper (Cu), ppm, (Min) 140

Selenium (Se), ppm, (Min) 0.6

Zinc (Zn), ppm (Min) 472

Vitamin A, I.U./lb, (Min) 20,000

Vitamin D, I.U./lb, (Min) 2,000

Vitamin E, I.U./lb, (Min) 10

INGREDIENTS**Note: ingredients differ by manufacturing plant**

Processed Grain By-Products, Calcium Carbonate, Forage Products, Urea, Salt, Calcium Chloride, Potassium Chloride, Soybean Oil, Thiamine Mononitrate, Vitamin A Supplement, Lecithin, Glyceryl Monostearate, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Zinc Sulfate, Colored with Iron Oxide, Basic Copper Chloride, Choline Chloride, Olive Oil, Vitamin D3 Supplement, Calcium Stearate, Roughage Products, Inositol, Sodium Selenite, Vitamin E Supplement.

MIXING & FEEDING DIRECTIONS:

Thoroughly mix 170 pounds of this Type B Medicated feed with 1830 pounds of good quality air dry coarse cracked or whole shelled corn and/or byproducts to obtain 1 ton of Type C medicated feed containing 25.5 g/ton Monensin.

Feed the resulting Type C Medicated feed continuously as sole ration at a rate of 19.25 pounds to provide 245 mg Monensin per head per day.

No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

CAUTION:

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions.

WARNING:

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Available Options:

Product No.	Package	Size Diameter	Feeding Rate
3003991-206	Bag	5/32 – 11/64	Follow Feeding Directions
3003991-201	Bulk	5/32 – 11/64	Follow Feeding Directions

Product Features:

Monensin

Contains protein, vitamins and minerals

Formulated using proprietary intake modifying technology

Product Benefits:

Supports the improvement in feed efficiency

Balances nutrient deficiencies in corn and by product-based ration to support growth.

Regulated number and size of meals which supports the digestive system in finishing cattle

Reduces the need for roughage in ration

Reduces storage and handling costs as well as decreasing manure production

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. This product should be fed with whole or coarse cracked corn. Cracked corn should be processed in a manner to minimize the creation of fines.
2. Whole shell corn should be at least of #2 corn quality.
3. Storage and handling of supplement should be managed to eliminate the unnecessary production of fines.
4. Do not allow animals to run out of feed.
5. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
6. When making a ration change, allow 7 to 10 days for animals to adjust to the new ration.
7. Provide adequate feeder space for each animal.
8. Provide mineral and salt free choice.
9. Provide fresh, clean water near the feeding area.

In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

KEY POINTS:

- The amount of ration consumed should be considered in determining the proper level of additive to be fed.
- All feeds containing moderate to high amounts of NPN can absorb moisture during periods of high relative humidity. The supplement may become hard in the sack or bridge in bulk bins. In extreme cases, pellets may become soft and crumble easily.