



SteakMaker® Finisher 40-20 R267

Specie: Beef**Type of Feed:** Supplement**Form of Feed:** Pellet**General Description:**

This economical supplement is designed for finishing cattle. The calcium level (7-8%) balances most grain based or silage-based diets with modest amounts of legume forages.

PURINA®

STEAKMAKER® FINISHER 40-20 R267**MEDICATED**

(For Ruminants Only)

SUPPLEMENT FEED FOR FINISHER FEEDLOT CATTLE

For improved feed efficiency in cattle fed in confinement for slaughter.

CAUTION: FEED AS DIRECTED**ACTIVE DRUG INGREDIENTS**

Monensin (as Monensin Sodium) 267.00 g/ton

GUARANTEED ANALYSIS

Crude Protein, (Min) 40.00 %
(This includes not more than 20% equivalent crude protein from non-protein nitrogen).

Crude Fat, (Min) 1.00 %

Crude Fiber, (Max) 12.00 %

Calcium (Ca), (Min) 7.00 %

Phosphorus (P), (Min) 0.50 %

Salt (NaCl), (Min) 4.00 %

Magnesium (Mg), (Min) 0.80%

Potassium (K), (Min) 1.00%

Copper (Cu), ppm, (Min) 200

Selenium (Se), ppm, (Min) 4.0

Zinc (Zn), ppm (Min) 700

Vitamin A, I.U./lb, (Min) 40,000

Vitamin D, I.U./lb, (Min) 10,000

Vitamin E, I.U./lb, (Min) 50

INGREDIENTS**Note: ingredients differ by manufacturing plant**

Processed Grain By-Products, Calcium Carbonate, Plant Protein Products, Urea, Animal Protein Products, Forage Products, Salt, Molasses Products, Magnesium Oxide, Calcium Lignin Sulfonate, Sodium Selenite, Vitamin D3 Supplement, Vitamin E Supplement, Vitamin A Supplement, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Zinc Sulfate, Basic Copper Chloride.

MIXING DIRECTIONS:

Thoroughly mix 150 pounds of this monensin Type B medicated feed with 1850 pounds of non-medicated feed to obtain one ton of Type C medicated feed containing 20 g/ton monensin.

FEEDING DIRECTIONS:

Feed the resulting Type C medicated feed continuously at a rate of 20 pounds to provide 200 mg of monensin per head per day.

WARNING:

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION:

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions.

Available Options:

Product No.	Package	Size Diameter	Feeding Rate
3004105-206	Bag	5/32-11/64	0.75 – 1.5 lb./hd./d
3004105-201	Bulk	5/32-11/64	0.75 – 1.5 lb./hd./d
3004105-101	Bulk	Meal	0.75 – 1.5 lb./hd./d
3004105-301	Crumble Bulk	5/32-11/64	0.75 – 1.5 lb./hd./d

Product Features:	Product Benefits:
Monensin	Supports the improvement in feed efficiency
40% Protein, 20 % maximum from NPN	Natural protein and modest level of NPN to meet protein requirements and optimize animal performance with economical cost of gain Product can be used in areas with higher humidity, yet has enough NPN to optimize cost of protein supplementation
7-8% Calcium, 0.5% Phosphorus	Balances moderately high-grain rations; helps reduce the likelihood of urinary calculi.
Adequate phosphorus for growing animals without being excessive	Optimizes animal performance while reducing environmental concerns with manure management
100% vitamin and trace mineral fortified	No need to feed extra mineral supplements to meet requirements

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Provide adequate bunk space for each animal. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
2. When making a ration change, allow 7-10 days for animals to adjust to the new ration.
3. Provide a source of fresh, clean water at all times.
4. Provide mineral and salt free choice.
5. When fed from a self-feeder, adjust feeder to minimize quantity of feed accumulating in trough.
6. Consult your veterinarian for recommended health program for your local area.
7. Have feed before cattle at all times. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

USAGE TIPS:

SteakMaker® 40-20 is a beef supplement for finishing cattle. It is ideal for use in moderate to high grain diets. It can fit well in these diets when fed with a wide range of roughage sources. It is typically fed at 0.75 to 1.5 lb. /hd. /d depending on the nutrient profile of other ration ingredients, ration energy content, age and size of cattle being fed and the medicated option used.

All ration adjustments should be made on a gradual basis. Use good bunk management techniques and procedures to prevent digestive disturbances and to optimize dry matter intake and animal performance. SteakMaker® 40-20 should be thoroughly mixed with grain and roughage. Do not top dress or feed separately. Follow label directions for mixing instructions.

KEY POINTS:

- The amount of ration consumed should be considered in determining the proper level of additive to be fed.
- All feeds containing moderate to high amounts of NPN can absorb moisture during periods of high relative humidity. The supplement may become hard in the sack or bridge in bulk bins. In extreme cases, pellets may become soft and crumble easily.