



SteakMaker® Grower 36-0 R200

Specie: Beef**Type of Feed:** Supplement**Form of Feed:** Pellet**General Description:**

This pelleted supplement is designed for growing and finishing cattle on moderately high grain diets.

PURINA®

STEAKMAKER® GROWER 36-0 R200

MEDICATED

(For Ruminants Only)

SUPPLEMENT FEED FOR GROWER FEEDLOT CATTLE

For improved feed efficiency in cattle fed in confinement for slaughter.

CAUTION: FEED AS DIRECTED**ACTIVE DRUG INGREDIENTS**

Monensin (as Monensin Sodium) 200.00 g/ton

GUARANTEED ANALYSIS

Crude Protein, (Min).....36.00 %
 Crude Fat, (Min).....1.00 %
 Crude Fiber, (Max)15.00 %
 Calcium (Ca), (Min)5.00 %
 Phosphorus (P), (Min).....0.80 %
 Salt (NaCl), (Min).....3.50 %
 Magnesium (Mg), (Min)0.50 %
 Potassium (K), (Min).....0.70 %
 Copper (Cu), ppm, (Min)135
 Selenium (Se), ppm, (Min)2.75
 Zinc (Zn), ppm (Min).....467
 Vitamin A, I.U./lb, (Min)27,000
 Vitamin D, I.U./lb, (Min)6,700
 Vitamin E, I.U./lb, (Min).....35

INGREDIENTSNote: ingredients differ by manufacturing plant

Grain Products, Roughage Products (not more than 40%), Calcium Carbonate, Processed Grain By-Products, Salt, Plant Protein Products, Molasses Products, Zinc Sulfate, Manganese Sulfate, Magnesium Oxide, Basic Copper Chloride, Sodium Selenite, Ethylenediamine Dihydriodide, Cobalt Carbonate, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement, Mineral Oil.

FEEDING DIRECTIONS:

Thoroughly mix 300 pounds of this monensin Type B medicated feed into 1700 pounds of non-medicated feed to obtain one ton of Type C medicated feed containing 30 g/ton monensin.

FEEDING DIRECTIONS:

Feed the resulting Type C medicated feed continuously at a rate of 16.7 pounds to provide 250 mg of monensin per head per day.

WARNING:

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION:

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions.

Available Options:

Product No.	Package	Size Diameter	Feeding Rate
3004152-201	Bulk	5/32-11/64	16.7 lb./hd./d (mixed as above)

Product Features:	Product Benefits:
Monensin	Improvement of feed efficiency in beef cattle
5% Calcium, 0.8% Phosphorus	Provides supplemental calcium to otherwise calcium deficient diets
Undegradable Intake Protein (UIP)	UIP supplementation supports nutrient demands for growth
100% vitamin and trace mineral fortified	No need to feed extra mineral supplements to meet requirements; fortified to optimize performance and support overall health
Pelleted	Prevents ingredient separation, helps assure you of consistent nutrient and feed additive delivery

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Do not allow animals to run out of feed.
2. Provide adequate bunk space for each animal. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed.
3. When making a ration change, allow 7-10 days for animals to adjust to the new ration.
4. Provide a source of fresh, clean water at all times.
5. Provide mineral and salt free choice.
6. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

USAGE TIPS:

SteakMaker® Grower supplement is pelleted, moderate calcium designed to be used in growing/finishing rations.

Recommended feeding rates of SteakMaker® Grower supplement is 1.5 pounds per head per day. Thoroughly mix SteakMaker® Grower supplement with the non-medicated ration prior to feeding.

KEY POINTS:

- Use in growing diets containing grains and a variety of forage sources
- Moderate feed rate may help lower cost of supplementation and can support cost of gain
- Provides proper level of vitamins, minerals and feed additives when fed according to label directions