



Spect*(i)*cle[®]
FLO

Product

information sheet

- // Active Ingredient: Indaziflam
- // Formulation: Flowable
- // WSSA Group: 29
- // Signal Word: No signal word required

Best Uses

- // Pre-emergent herbicide controls the most troublesome grassy and broadleaf weeds, as well as annual sedges
- // Extended residual control of annual bluegrass (*Poa annua*), goosegrass, crabgrass and 75 more weeds

Attributes of Specticle[®] FLO

- // Controls weeds by inhibiting cellulose biosynthesis – its primary site of action is the newly developing root of a germinated weed seedling
- // No stains, no odor formulation
- // Mixable with post-emergent herbicides
- // Newer mode-of-action – controls weeds known to develop resistance to other herbicides
- // Works in turf, landscapes and hardscapes

How to Apply

- // Apply to turf before weed seeds germinate
- // Use as a broadcast treatment or directed spray on bare ground or mulched areas around labeled ornamental plants and grasses
- // Adequate soil moisture is needed to activate Specticle FLO and provide maximum control
- // If weeds are present, tank-mix Specticle FLO with selective, post-emergent herbicides like Revolver[®], Celsius[®] WG or Tribute[®] Total
- // To control emerged weeds in plant beds, dormant turf, cart paths or parking lots, use in conjunction with a non-selective herbicide



Poa annua



Eleusine indica



Digitaria sanguinalis

Key Strengths

- // Broad-spectrum, preemergence herbicide
- // Long residual
- // Low use rates
- // Broadleaf and grassy weed control

Use Rates for Single Application of Specticle FLO**

Target Weed (single application)		Specticle FLO Use Rate (fl oz per acre)
Crabgrass Goosegrass	Annual bluegrass Broadleaf weeds	6.0 - 10.0
Annual sedge	Kyllinga	10.0

Use Rates for Split Application of Specticle FLO**

Target Weed	Initial Application Use Rate (fl oz per acre)	Follow-up Application (within 45 - 90 days) Use Rate (fl oz per acre)
Crabgrass Goosegrass Annual bluegrass	4.5 - 9.0	3.0 - 9.0
Broadleaf weeds	6.0 - 9.0	3.0 - 9.0
Annual sedge Kyllinga	9.0	6.0 - 9.0

Use Rates for Multiple Applications of Specticle FLO**

Target Weed	Initial Application Use Rate (fl oz per acre)	Second Application (within 30 - 45 days) Use Rate (fl oz per acre)	Third Application (within 30 - 45 days) Use Rate (fl oz per acre)
Crabgrass Goosegrass Annual bluegrass	3.0 - 4.5	3.0 - 4.5	3.0 - 4.5

*Specticle FLO 2(ee) allows these use rates on annual bluegrass in the following states: AL, AZ, AR, CO, FL, GA, HI, IN, IL, KS, LA, MD, MS, MO, NC, OH, OK, SC, TN, TX, UT, VA, WV.

**Do not exceed 18.5 fl. oz./A of Specticle FLO for all applications within a 12-month period.



Goosegrass

Large crabgrass

Spotted spurge

Doveweed

www.EnvironmentalScience.Bayer.US



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS
Bayer Environmental Science, a Division of Bayer CropScience LP, 5000 CentreGreen Way, Suite 400, Cary, NC 27513. For additional product information, call toll-free 1-800-331-2867. Not all products are registered in all states. Bayer, the Bayer Cross, Specticle, Celsius, Tribute and Revolver are registered trademarks of Bayer. © 2019 Bayer CropScience LP. ES-818-SPECFLO-167-A-R2