



Specticle® G

Product information sheet

- // Active Ingredient: Indaziflam
- // Formulation: Granule
- // WSSA Group: 29
- // Signal Word: No signal word required

Best Uses

- // Granular preemergence herbicide controls the most troublesome grassy and broadleaf weeds, as well as annual sedges
- // Extended residual control of annual bluegrass (*Poa annua*), goosegrass, crabgrass and more than 90 listed weeds

Attributes of Specticle® G

- // Controls weeds by inhibiting cellulose biosynthesis – its primary site of action is the newly developing root of a germinated weed seedling
- // No stains, no odor formulation
- // Newer mode-of-action – controls weeds known to develop resistance to other herbicides
- // Works in turf, landscapes and hardscapes

How to Apply

- // Apply before weed seeds germinate
- // Use as a broadcast treatment on bare ground or mulched areas around labeled ornamental plants and grasses
- // Specticle G performs best when irrigation or rainfall occurs prior to weed seed germination
- // Adequate soil moisture is needed to activate Specticle G and provide maximum control



Poa annua

Eleusine indica

Digitaria sanguinalis

Key Strengths

- // Broad-spectrum, preemergence herbicide
- // Long residual
- // Low use rates
- // Broadleaf and grassy weed control

Use Rates For Single Application Of Specticle G

Target Weed*		Specticle G Use Rate (lbs per acre)**	Specticle G Use Rate (lbs per 1,000 sq ft)
Crabgrass Goosegrass	Annual bluegrass Broadleaf weeds	100 - 200	2.3 - 4.6
Annual sedge	Kyllinga	200	4.6

Use Rates For Split Application Of Specticle G


Target Weed*		Specticle G Use Rate (initial application)		Specticle G Use Rate (within 90 days of initial application)	
		(lbs per acre)**	(lbs per 1,000 sq ft)	(lbs per acre)**	(lbs per 1,000 sq ft)
Crabgrass Goosegrass	Annual bluegrass Broadleaf weeds	100 - 200	2.3 - 4.6	100 - 200	2.3 - 4.6
Annual sedge	Kyllinga	200	4.6	125 - 200	2.9 - 4.6

*Use higher use rates within the rate range for each target weed when a site has historically higher weed pressure.

**The maximum single application on turf is 200 lbs./A (4.6 lbs./1,000 sq. ft.). Do not exceed 400 lbs. of Specticle G/A (9.2 lbs./1,000 sq. ft.) in a 12-month period.



Goosegrass (Dr. Bert McCarty,
Clemson University)



Large Crabgrass (Dr. Bert McCarty,
Clemson University)



Spotted Spurge (Dr. Bert McCarty,
Clemson University)



Florida Pusley (Dr. Bert McCarty,
Clemson University)



Yellow Woodsorrel (Dr. Bert McCarty,
Clemson University)

www.EnvironmentalScience.Bayer.US



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS
Bayer Environmental Science, a Division of Bayer CropScience LP, 5000 CentreGreen Way, Suite 400, Cary, NC 27513. For additional product information, call toll-free 1-800-331-2867. Not all products are registered in all states. Bayer, the Bayer Cross and Specticle are registered trademarks of Bayer.
© 2019 Bayer CropScience LP. ES-818-SPECG-168-A-R2